

Vacuum stress around a topological defect

V. A. De Lorenci^{a*} and E. S. Moreira, Jr.^{a†}

^aInstituto de Ciências, Universidade Federal de Itajubá,
Av. BPS 1303 Pinheirinho, 37500-903 Itajubá, MG, Brazil

We show that a dispiration (a disclination plus a screw dislocation) polarizes the vacuum of a scalar field giving rise to an energy momentum tensor which, as seen from a local inertial frame, presents non vanishing off-diagonal components. The results may have applications in cosmology (chiral cosmic strings) and condensed matter physics (materials with linear defects).

It is fairly well known that a needle solenoid carrying a magnetic flux makes virtual charged particles to run around the solenoid inducing a non vanishing current density (see e.g. Ref [1]). We wish to consider what seems to be a gravitational (geometric) analogue of this Aharonov-Bohm effect, by computing the vacuum expectation value of the energy momentum tensor of a massless and neutral scalar field far away from a dispiration.

Let us begin by presenting the geometry of the background (units are such that $c = \hbar = 1$),

$$ds^2 = dt^2 - dr^2 - \alpha^2 r^2 d\theta^2 - (dz + \kappa d\theta)^2, \quad (1)$$

where the points labeled by (t, r, θ, z) and $(t, r, \theta + 2\pi, z)$ are identified [2,3]. When $\alpha = 1$ and $\kappa = 0$ Eq (1) becomes the line element of the flat space-time written in cylindrical coordinates. Borrowing terminologies in condensed matter physics, the parameters α and κ correspond to a disclination and a screw dislocation, respectively. We should remark that Eq (1) may be associated with the gravitational background of certain chiral cosmic strings [4] (as has been suggested in Ref. [2]), as well as can describe (in the continuum limit) the effective geometry around a dispiration in an elastic solid (see Ref. [5] and references therein).

The definitions $\varphi := \alpha\theta$ and $Z := z + \kappa\theta$ lead to

$$ds^2 = dt^2 - dr^2 - r^2 d\varphi^2 - dZ^2, \quad (2)$$

which should be considered together with the peculiar identification

$$(t, r, \varphi, Z) \sim (t, r, \varphi + 2\pi\alpha, Z + 2\pi\kappa). \quad (3)$$

Although Eq. (2) expresses the fact that the background is locally flat, due to Eq. (3) we cannot use Eq. (2) (which is a local statement) to infer that the global symmetries of the background are the same as those of the Minkowski space-time (in this sense Eq. (2) is singular). In fact, Eq. (2) disguises a curvature singularity on the symmetry axis [2] (when $\kappa \neq 0$, in the context of the Einstein-Cartan theory, there is also a torsion singularity at $r = 0$ [3,6]).

The vacuum expectation value of the energy momentum tensor is obtained by applying a differential operator to the renormalized scalar propagator around a dispiration (see e.g. Ref. [7]),

$$\langle T^\mu{}_\nu \rangle = i \lim_{x' \rightarrow x} \mathcal{D}^\mu{}_\nu(x, x') D^{(\alpha, \kappa)}(x, x'). \quad (4)$$

We have recently obtained $D^{(\alpha, \kappa)}(x, x')$ (classical propagators have been considered in Ref [8]) by using the Schwinger proper time prescription combined with the completeness relation of the eigenfunctions of the d'Alembertian operator [9]. Such eigenfunctions have the form $R(r)\chi(\varphi)\exp\{i(\nu Z - \omega t)\}$ which, by observing Eq. (3), leads to

$$\chi(\varphi + 2\pi\alpha) = e^{-i2\pi\nu\kappa}\chi(\varphi). \quad (5)$$

This boundary condition is typical of the Aharonov-Bohm set up where $\nu\kappa$ is identified

*delorenci@unifei.edu.br

†moreira@unifei.edu.br

with the flux parameter $e\Phi/2\pi$. If we carry over to the four-dimensional context lessons from gravity in three dimensions [10,11], it follows that the charge e and the magnetic flux Φ should be identified with the longitudinal linear momentum ν and $2\pi\kappa$, respectively [2].

When $\kappa/r \rightarrow 0$, Eq. (4) yields for the diagonal components the expressions of the vacuum fluctuations around an ordinary cosmic string ($\kappa = 0$) [12]. Regarding the other components, the prescription in Eq. (4) kills off the dominant contribution in the renormalized propagator [9], resulting that the subleading contribution yields two non vanishing off-diagonal components,

$$\langle T^\varphi_Z \rangle = \frac{i}{r^2} \lim_{x' \rightarrow x} \partial_\varphi \partial_Z D^{(\alpha, \kappa)}(x, x') = \frac{\kappa}{r^6} B(\alpha), \quad (6)$$

and

$$\langle T^Z_\varphi \rangle = \frac{\kappa}{r^4} B(\alpha), \quad (7)$$

where $B(\alpha)$ depends on the disclination parameter only [9]. Unlike the diagonal components, $\langle T^\varphi_Z \rangle$ and $\langle T^Z_\varphi \rangle$ do not depend on the coupling parameter ξ .

When $\alpha = 1$, $B = 1/60\pi^2$ which corresponds approximately to the value of α in the physics of formation of ordinary cosmic strings [13].

It is instructive to display both disclination and screw dislocation effects in a same array. When $\xi = 1/6$ (conformal coupling), for example, $\langle T^\mu_\nu \rangle$ with respect to the local inertial frame [cf. Eq. (2)] can be cast into the form

$$\langle T^\mu_\nu \rangle = \frac{1}{r^4} \begin{pmatrix} -A & 0 & 0 & 0 \\ 0 & -A & 0 & 0 \\ 0 & 0 & 3A & \kappa B/r^2 \\ 0 & 0 & \kappa B & -A \end{pmatrix}, \quad (8)$$

where $A(\alpha) := (\alpha^{-4} - 1)/1440\pi^2$, and which holds far away from the defect (and for $\alpha \neq 1$, when $\kappa \neq 0$). [When $\kappa \neq 0$, by setting $\alpha = 1$ in Eq. (8), A vanishes and subleading contributions depending on κ take over.]

Before closing we should report a discrepancy in the literature. Recently, works [14,15] have appeared stating that, for $\alpha = 1$, the subleading contribution in $\langle T^0_0 \rangle$ [cf. Eq. (8)] is $\lambda\kappa^2/r^6$, where λ is a certain constant. We have

performed a rather detailed study showing that such a subleading contribution is $f(\kappa/r)\kappa^2/r^6$ instead, where $f(\kappa/r)$ diverges as $\kappa \rightarrow 0$ [although $\lim_{\kappa \rightarrow 0} \kappa^2 f(\kappa/r) = 0$].

acknowledgments

This work was partially supported by the Brazilian research agencies CNPq and FAPEMIG.

REFERENCES

1. E.M. Serebryanyi, *Theor. Math. Phys.* 64 (1985) 846.
2. D.V. Gal'tsov and P.S. Letelier, *Phys. Rev. D* 47 (1993) 4273.
3. K.P. Tod, *Class. Quant. Grav.* 11 (1994) 1331.
4. J.D. Bekenstein, *Phys. Rev. D* 45 (1992) 2794.
5. R.A. Puntigam and H.H. Soleng, *Class. Quant. Grav.* 14 (1997) 1129.
6. P.S. Letelier, *Class. Quant. Grav.* 12 (1995) 471.
7. N.D. Birrel and P.C.W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, Cambridge, England, 1982.
8. V.A. De Lorenci and E.S. Moreira, Jr., *Phys. Rev. D* 65 (2002) 107503.
9. V.A. De Lorenci and E.S. Moreira Jr., hep-th/0301194.
10. P.S. Gerbert and R. Jackiw, *Commun. Math. Phys.* 124 (1989) 229.
11. P.S. Gerbert, *Nucl. Phys. B* 346 (1990) 440.
12. M.E.X. Guimarães and B. Linet, *Commun. Math. Phys.* 165 (1994) 297.
13. A. Vilenkin and E.P.S. Shellard, *Cosmic Strings and Other Topological Defects*, Cambridge University Press, Cambridge, England, 1994.
14. I. Pontual and F. Moraes, *Philos. Mag. A* 78 (1998) 1073.
15. T.M. Helliwell, D.A. Konkowski, and V. Arndt, *Gen. Rel. Grav.* 35 (2003) 79.